

Meeting abstract

Unexpected role of STAT1 serine727 for NK cell function

Eva M Putz¹, Eva Zebedin¹, Agnes Csiszar², Michaela Prchal³,
Angelika Berger¹, Thomas Decker⁴, Pavel Kovarik⁴ and Veronika Sexl^{*1}

Address: ¹Institute of Pharmacology, Center for Biomolecular Medicine and Pharmacology, Medical University of Vienna, 1090 Vienna, Austria, ²Research Institute of Molecular Pathology, 1030 Vienna, Austria, ³Institute of Animal Breeding and Genetics, University of Veterinary Medicine, 1210 Vienna, Austria and ⁴Max F. Perutz Laboratories, University of Vienna, 1030 Vienna, Austria

Email: Veronika Sexl* - veronika.sexl@meduniwien.ac.at

* Corresponding author

from 15th Scientific Symposium of the Austrian Pharmacological Society (APHAR) Joint meeting with the Hungarian Society of Experimental and Clinical Pharmacology (MFT) and the Slovenian Pharmacological Society (SDF)
Graz, Austria. 19-21 November 2009

Published: 12 November 2009

BMC Pharmacology 2009, 9(Suppl 2):A29 doi:10.1186/1471-2210-9-S2-A29

This abstract is available from: <http://www.biomedcentral.com/1471-2210/9/S2/A29>

© 2009 Putz et al; licensee BioMed Central Ltd.

Background

Natural killer (NK) cells are important key players of the innate immune system and provide immediate defense against viral infection and tumor transformation. Therefore, their potential in cancer immunotherapy has grown prominently in the last years [1]. As is known from the literature, the Signal Transducer and Activator of Transcription 1 (STAT1) plays an important role for NK cell function, since STAT1-deficient mice display impaired basal NK cytolytic activity *in vitro* and are unable to reject transplanted tumors *in vivo* [2]. STAT1 mediates signals downstream of interferons and gets activated by phosphorylation of several tyrosine and serine residues. In particular phosphorylation of S727 is considered a prerequisite for the full-fledged activation of STAT1 [3]. The aim of this project is to investigate the role of STAT1-S727 in NK cell-mediated cytotoxicity and tumor surveillance.

Methods

The analysis of STAT1^{-/-}, STAT1-S727A and wild-type mice includes the preparation of primary splenic NK cells and their investigation regarding proliferation ([³H]thymidine-incorporation), cytotoxicity (standard [⁵¹Cr]-release) and cytokine production (ELISA, multi-plex bead arrays). *In vivo* tumor models are employed using NK-sensitive tumor cell lines (v-abl⁺ leukemia, B16 melanoma, 4T1 breast cancer).

Results

Wild-type NK cells from untreated healthy mice display basal phosphorylation on STAT1-S727 *in vivo*. Surprisingly, disruption of this phosphorylation site by mutating serine727 to alanine (STAT1-S727A) significantly enhances NK cell cytotoxicity towards various target cells *in vitro* compared to wild-type. Moreover, we demonstrate that STAT1-S727A mice do not only display delayed leukemia onset but also lower susceptibility to intravenously administered B16 (melanoma) and 4T1 (metastasizing breast cancer) cells compared to STAT1^{-/-} and wild-type mice.

Conclusion

Obviating the phosphorylation of STAT1-S727 seems to increase NK cell cytotoxicity *in vitro* and *in vivo*. As an ultimate goal we aim to find the underlying mechanism(s) and upstream regulator(s), since loosening the brake on NK cell functions could represent a potential novel strategy in cancer therapy.

Acknowledgements

We thank the Austrian Academy of Science for supporting Eva Maria Putz with a DOC-ffORTE fellowship.

References

1. Sutlu T, Alici E: **Natural killer cell-based immunotherapy in cancer: current insights and future prospects.** *J Intern Med* 2009, **266**:154-181.

2. Lee CK, Rao DT, Gertner R, Gimeno R, Frey AB, Levy DE: **Distinct requirements for IFNs and STAT1 in NK cell function.** *J Immunol* 2000, **165**:3571-3577.
3. Varinou L, Ramsauer K, Karaghiosoff M, Kolbe T, Pfeffer K, Muller M, Decker T: **Phosphorylation of the Stat1 transactivation domain is required for full-fledged IFN- γ -dependent innate immunity.** *Immunity* 2003, **19**:793-802.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

